

Climate Change, Flood Disaster Assessment and Human Security in Katsina State, Nigeria

I.B. Abaje¹, A.O. Ogoh², B.B. Amos³ and M. Abashiya³

¹*Department of Geography & Regional Planning, Federal University, Dutsin-Ma, Katsina State*

²*Department of Political Science, Federal University, Dutsin-Ma, Katsina State*

³*Department of Geography, Gombe State University, Gombe*

This study examines the recent rainfall trends and occurrence of floods in Katsina State. Annual and monthly rainfall data for Katsina from the period 1985 to 2014 were used to analyze the rainfall trends. The standard deviation was then used to determine the rainfall deviation from the normal. The paper focused on the floods event of 2012 and 2013 in the state and the implications for human security. Floods data were obtained from Katsina State Emergency Management Agency, and a total of 200 copies of a questionnaire were administered to flood affected communities in order to obtain information of the causes and damages from the flood hazards of 2012 and 2013. Trend analysis was used to determine the rainfall changes and the results show an increase in rainfall amount in the months of June, July and August which is one of the factors responsible for the frequent occurrences of floods in those months. The major implication of this finding is that infrastructures built on the perceived decreasing rainfall have to be reviewed. The study recommends that, for sustainable development, planning and designing of infrastructures should take into account the recent rainfall trends and occurrences of floods in the area.

Key Words: climate change, floods, hazard, human security, infrastructures, trends

Introduction

Climate change is one of the greatest socioeconomic and biophysical challenges confronting the world in the 21st century. Human activity, particularly deforestation and the burning of fossil fuels, is driving this change by increasing atmospheric concentrations of carbon dioxide and other greenhouse gases (GHGs). As a result, the world is experiencing greater weather extremes, changes in rainfall patterns, heat and cold waves, and increasing droughts and floods. These phenomena have a negative impact on the environment and on people's lives and livelihoods. Marginalized groups and communities in the poorest regions of the world are particularly affected, even as they are least responsible for these changes (United Nations Development Programme [UNDP], 2009).

Flood disaster which are always the result of both natural phenomena and human actions has increased in number and magnitude in many regions of the world over this few years thereby demanding attention and actions. According to UNDP (2004) flooding is the outcome of natural hazard and human vulnerability coming together. Indeed, flood disaster do not only happen in a physical environment, it also occur in a social and political context, this imply that

flood disaster not only reveal the underlying social, economic, political and environment problems, but unfortunately they contribute to worsening them, hindering economic and social progress. There is increasing evidence that pre-existing social, economic, political and environmental conditions determine the impact of flood disaster, thus, the weaker the pre-existing conditions are, the higher the impact of a flood disaster will be. In other words, the more vulnerable, and thus less resilient, are the social context in which a disaster occurs, the bigger the impact and the damage. As the flood disaster occurred, the pre-existing structure and social condition actually determined the impact it had on the people and social structure.

Corresponding author: I B Abaje, Department of Geography and Regional Planning, Federal University, Dutsin-Ma, P.M.B. 5001, Dutsin-Ma, Katsina State, Nigeria. E-mail: abajebest@gmail.com

 This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

The severity of the impacts of the flood disaster is seen in the displacement of over 21million people; destruction of 597,476 houses; death of 363 people and estimated loss of USD 19.6 billion in the country (Nigeria Emergency Management Agency [NEMA], 2013), thereby raising security issues as it broaden security concerns away from exclusive military threats to state to include non-military threats like poverty, hunger, diseases, environmental degradation, thus the concept of human security.

Traditionally, flood disaster is conceived as environmental or humanitarian issue, but they now constitute security concerns. So in today's world, and because of their probable relationship to climate change, flood disaster must essentially be understood as a human security issue. They represent not only a threat to people's survival, but also constitute a vulnerability factor for people and communities and even nations.

Flooding as witnessed in some parts of Katsina state in the years under study (2012 and 2013) severely impacted the human security of areas hit by the disaster. On an individual or household level, the flood brought about death, injuries and trauma. The floods destroyed people's livelihoods by massively devastating farmland, farm produce and social infrastructures, thereby becoming dependent on relief assistance from government, NGOs and other organization to survive.

In Katsina State, limited studies have been conducted on flood disaster and human security. The specific objectives of this study are to: 1) examine the rainfall characteristics of the study area; 2) examine the causes of floods in the area with particular reference to the 2012 and 2013 flood disasters; and 3) examine the damages and impacts on the people of the study area

Study Area

Katsina State (Figure 1) is located between latitude 11⁰00'N and 13⁰20'N and longitude 7⁰00'E and

8⁰55'E. It shares boarder with Niger Republic to the North, Kaduna State to the South, Jigawa and Kano States to the East, and Zamfara State to the West. Katsina State has a land size of about 24,971.215km² with a population of 5,801,584 as at 2006 national census (Federal Republic of Nigeria, 2012).

The climate of Katsina State is the tropical wet and dry type (Tropical Continental Climate), classified by Koppen as Aw climate. Rainfall is between May and September with very high intensity between the months of July and August (Abaje, Sawa & Ati, 2014). The average annual rainfall varies from 550 mm in the northern part to about 1000 mm in the southern part of the state. The pattern of rainfall in the state is highly variable. As a result, the state is subject to frequent floods that can impose serious socio-economic constraints (Abaje, Ati & Iguisi, 2012a).

Seasonal variation in rainfall is directly influenced by the interaction of two air masses: the relative warm and moist tropical maritime (mT) air mass, which originates from the Atlantic Ocean associated with southwest winds in Nigeria; and the relatively cool, dry and stable tropical continental (cT) air mass that originates from the Sahara Desert and is associated with the dry, cool and dusty North-East Trades known as the Harmattan. The boundary zone between these two air streams is called the Inter-tropical Discontinuity (ITD). The movement of the ITD northwards across the state in August marks the height of the rainy season in the whole state while its movement to the southernmost part around February marks the peak of the dry season in the state (Odekunle, 2006; Abaje *et al*, 2012b; Abaje *et al*, 2014). The annual mean temperature is about 27⁰C. The highest air temperature normally occurs in April/May and the lowest in December through February (Nigerian Meteorological Agency, 2012). The soil is sandy in nature while the vegetation is the Sudan Savanna type which combines the characteristics and species of both the Guinea and Sahel Savanna (Abaje, 2007).




Figure 1: Map of Katsina State Showing the Study Area

Materials and Methods

Rainfall data for thirty (30) years (1985-2014) were used to analyze the recent rainfall characteristics of the study area. The data were sourced from the archive of Nigerian Meteorological Agency (NIMET). Only rainfall totals for the months of May to September and the annual totals were used in this study. The main reason is that 85% of the total annual rainfall is

received in the study area are within these months (May to September).

The standardized coefficients of Skewness (Z_1) and Kurtosis (Z_2) statistics as defined by Brazel and Balling (1986) were used to test for the normality in the rainfall series for the study area. The standardized coefficient of Skewness (Z_1) was calculated as:

$$Z_1 = \left[\frac{\left(\sum_{i=1}^N (x_i - \bar{x})^3 / N \right)}{\left(\sum_{i=1}^N (x_i - \bar{x})^2 / N \right)^{3/2}} \right] / \left(\frac{6}{N} \right)^{1/2}$$

and the standardized coefficient of Kurtosis (Z_2) was determined as:

$$Z_2 = \left[\frac{\left(\sum_{i=1}^N (x_i - \bar{x})^4 / N \right)}{\left(\sum_{i=1}^N (x_i - \bar{x})^2 / N \right)^2} \right] - 3 / \left(\frac{24}{N} \right)^{1/2}$$

where \bar{x} is the long term mean of x_i values, and N is the number of years in the sample. These statistics were used to test the null hypothesis that the individual temporal samples came from a population with a normal (Gaussian) distribution. If the absolute value of Z_1 or Z_2 is greater than 1.96, a significant deviation from the normal curve is indicated at the 95% confidence level. If the data are not found to be normally distributed, various transformation models

could be used to normalize the series such as Log transformation and Lambda transformations of Box and Cox (1964), and Square and Cube Root transformations (Stidd, 1970) amongst others.

The Relative Seasonality Index (SI) of the rainfall series was calculated using the Walsh and Lawler (1981) statistic. This was done in order to show the class into which the climate of Katsina State can be classified. This index is calculated as:

$$SI = \frac{1}{R} \sum_{n=1}^{n=12} \left| \bar{x}_n - \frac{\bar{R}}{12} \right| \text{----- (1)}$$

where \bar{x}_n is the mean rainfall for month n and \bar{R} is the mean annual rainfall. This index can vary from zero (if all the months have equal rainfall) to 1.83 (if

all the rainfall occurs in a single month). Table 1 shows the seasonality index classes as proposed by Walsh and Lawler (1981).

Table 1: Seasonality index classes

| Rainfall Regime | SI Class Limits |
|--|-----------------|
| Very equable | ≤ 0.19 |
| Equable but with a definite wetter season | 0.20-0.39 |
| Rather seasonal with a short drier season | 0.40-0.59 |
| Seasonal | 0.60-0.79 |
| Markedly seasonal with a long drier season | 0.80-0.99 |
| Most rain in 3 months or less | 1.00-1.19 |
| Extreme, almost all rain in 1-2 months | ≥ 1.20 |

To examine the nature of the trend, linear trend lines and moving mean were calculated and plotted using Microsoft Excel statistical tool (2013) for both the monthly (May to September) and annual rainfall totals (mm) for the station. In this work, the 5-year moving mean was used in order to smoothing the time series, thereby reducing the irregular fluctua-

tions and highlighting those that are regular. The standard deviation which has the potential to provide a result of deviation from normal was equally determined and plotted using Microsoft Excel statistical tool (2013). The standard deviation (δ) is given by the formula:

$$\delta = \sqrt{\sum \frac{(X - \bar{X})^2}{N}} \text{----- (2)}$$

A total of 200 copies of a questionnaire were administered to flood victims of 2012 and 2013 in the affected Local Government Areas (LGAs) of the state. This was done in order to obtain information about their perceptual assessment of the causes and damages from the flood disaster. Eighteen (18) LGAs were affected in the 2012 flood disaster while the 2013 flood disaster affected twenty (20) LGAs (Table 2). The 2012 and 2013 floods affected LGAs were then ranked in descending order based on the affected number of victims according to their senatorial districts. In a situation where a LGA is affected in both the 2012 and 2013 flood disasters, the year with the highest number of victims was then used in the rank-

ing. Proportional sampling was used in selecting the sampled LGAs in the three senatorial districts, Five LGAs with the highest number of victims were purposely selected as follows: Dutsi and Sandamu LGAs in Daura Senatorial District, Batagarawa and Katsina LGAs in Katsina Senatorial District, and Matazu LGA in Funtua Senatorial District. The flood victims were then administered questionnaires based on simple proportion (Table 3). These LGAs were also chosen as an attempt to provide a representative sample of the victims' perceptual assessment of the causes and damages from the 2012 and 2013 flood disasters within the constraints of time and available resources.

Table 2: Flood Disaster Affected Local Government Areas (2012 and 2013)

| 2012 Flood Disaster | | 2013 Flood Disaster | |
|---------------------|------------------|---------------------|------------------|
| LGA | Affected Victims | LGA | Affected Victims |
| Maiadua | 721 | Charanchi | 129 |
| Batsari | 409 | Kankia | 225 |
| Bindawa | 514 | Daura | 281 |
| Safana | 475 | Zango | 257 |
| Kurfi | 244 | Mashi | 81 |
| Kankia | 23 | Katsina | 285 |
| Jibia | 17 | Malumfashi | 102 |
| Katsina | 535 | Ingawa | 32 |
| Dutsi | 949 | Musawa | 223 |
| Ingawa | 785 | Matazu | 153 |
| Rimi | 425 | Danja | 90 |
| Mani | 489 | Dutsin-Ma | 93 |
| Daura | 608 | Jibia | 241 |
| Matazu | 1365 | Batagarawa | 331 |
| Sandamu | 948 | Kaita | 279 |
| Batagarawa | 1232 | Funtua | 227 |
| Danmusa | 259 | Baure | 53 |
| Kusada | 515 | Kurfi | 186 |
| | | Maiadua | 281 |
| | | Danmusa | 94 |

Source: Katsina State Emergency Management Agency (2014)

Key informant survey was also carried out with field staff of the State Emergency Management Agency

(SEMA) in order to obtain in-depth information on the causes and damages of the 2012 and 2013 floods.

Table 3: Selected Local Government Area in Each Senatorial District

| Senatorial District | LGAs | Selected LGAs | No of Affected Victims | No. Sampled |
|---------------------|--|---------------|------------------------|-------------|
| Daura | Dutsi, Sandamu, Ingawa, Maiadua, Daura, Kusada, Bindawa, Mani, Zango, Kankia, Mashi, and Baure. | Dutsi | 949 | 38 |
| | | Sandamu | 948 | 38 |
| Katsina | Batagarawa, Katsina, Safana, Rimi, Batsari, Kaita, Danmusa, Kurfi, Jibia, Charanchi, and Dutsin-Ma | Batagarawa | 1232 | 49 |
| | | Katsina | 535 | 21 |
| Funtua | Matazu, Funtua, Musawa, Malumfashi, and Danja | Matazu | 1365 | 54 |
| Total | | | | 200 |

Source: Field Survey, 2014

Results and Discussion

Rainfall Characteristics

The calculated SI was 1.13, which means that most of the rainfall in the study area is within 3 months or less. The general statistics of the monthly (May to September) and annual rainfall of the study area


(1985-2014) are presented in Table 4. The results of Z_1 and Z_2 show that all the months and the annual were accepted as indicative of normality at the 95% significant level. Based on these results, no transformation was done to the data. The mean monthly rainfall of the study period shows a single peak in August (194.8 mm).

Table 4: General statistics of the rainfall in mm (1985-2014)

| Statistics | May | June | July | August | September | Annual |
|------------|-------|-------|-------|--------|-----------|--------|
| \bar{X} | 34.6 | 74.2 | 134.6 | 194.8 | 88.4 | 544.7 |
| SD | 31.5 | 43.4 | 52.7 | 77.7 | 53.1 | 170.8 |
| Z_1 | 0.63 | 0.64 | 0.54 | 0.41 | 1.15 | 0.02 |
| Z_2 | -1.03 | -0.88 | -0.09 | -0.93 | 0.62 | -0.23 |

Figure 2 (a) shows the graphical presentation of the annual trends of the rainfall series smoothed out with the 5-year running mean. The lowest annual rainfall of about 240.6 mm was experienced in 1985 while the highest annual rainfall of about 955.7 mm was experienced in 2010. The 5-year running mean shows annual rainfall below the long-term mean last-

ed from the beginning of the data up to the late 1990s. From the late 1990s to the end of the study period, the rainfall was above the long-term mean. The linear trend line also show an increase in rainfall. A general examination of the monthly running means (Figure 2 (b-f)) shows that rainfall has been increasing from the year 2000 to the end of the study period.






Figure 2: Rainfall Trends for Annual and Monthly Totals





Figure 2: Continued

The plotted standard deviation for the monthly rainfall anomalies (Figure 4b-f) generally show that years of rainfall above the mean standard deviation were more than below the mean standard deviation from the late 1990s to the end of the study period. The annual rainfall (Figure 4a) shows 9 anomaly years (5 below the mean standard deviation and 4 above it) of the 30 years of study. All the 5 years of rainfall below the mean standard deviation occurred between 1985 and 1993. These years of rainfall below the mean standard deviation coincides with the droughts

of the 1980s that ravaged the country. On the other hand, all the 4 years of rainfall above the mean standard deviation occurred between the 2005 and 2013. The findings is also in good agreement with the observations made by Odekunle, Andrew and Aremu (2008), Abaje *et al* (2012a&b), Abaje, Ati, Iguisi, and Jidauna (2013) and Abaje, Ndabula and Garba (2014) that northern Nigeria has been experiencing decreasing number of dry conditions and consequently, increasing wetness from the 1990s to the recent years. The increasing wetness appears to be accounted for

by significant northward shifts in the surface location of the ITD over the country.

It is clear from the results that the annual rainfall has increase in the last 30 years. Increase in the annual rainfall yield is predominantly as a result of the increase in June, July and August rainfall. The increase in rainfall for these months is one of the factors responsible for the frequent occurrence of floods within those months.

Socioeconomic Characteristics of Respondents

The results of the field survey revealed that the majority of the respondents were males (89.5%) while only 10.5% were females. 82.5% of the respondents are married; and the average household size is 9. Out of the 200 respondents, 23.5% attended Quranic School, 34% primary school, 12% have tertiary education, and 24.5% have secondary education, while 6.0% have no education at all. Majority of the respondents (43.5%) are between 41-50 years. The major occupation of the respondents is farming which represent 55.5%, while 24.5% are civil servants, 16.5% traders, and others (3.5%). This implies that, the respondents depend heavily on natural and physical resources of the environment for their livelihood.

Causes of Floods

About 54.5% of the respondents opined that rainfall has been increasing in recent times. This is in agreement with the analyzed rainfall data that showed an increase in rainfall over the recent years. The result is also in agreement with the study of Abaje *et al* (2012b) in the Sudano-Sahelian Zone of Nigeria that,

at present, the climate of the region indicates a tendency towards a wetter condition rather than the increasing dryness that was a feature of the period from the 1960s to the 1980s. Heavy rainstorms according to the respondents (63.5%) do precede the occurrence of floods in the area and was suggested as the main cause of the 2012 and 2013 floods. This result is in agreement with the work of Abaje and Giwa (2010) in which 87% of the respondents said that torrential rainstorm do precede the occurrence of floods. About 65.5% of the respondents said that August rainfall is always heavy and the month marks the peak of the raining season in the study area. Only 19.5% and 15% opined that the rainfall is heavy in the months of July and September respectively.

In 2012, the highest amount of rainfall was recorded in the month of July (224 mm) while August recorded 195.8 mm that year; whereas in 2013, the highest amount of rainfall (284.7 mm) was recorded in the month of August (Figure 3). It means that rainfall amount in the state is concentrated within two months (July and August). Coincidentally, the 2012 flood disaster occurred in the months of July and August while the 2013 flood disaster occurred in the month of August. This means that the heavy rainstorm of the months of July and August is one of the factors responsible for the occurrence of the July 2012 and August 2013 floods in the area. Climate change may be responsible for the unusual and frequent torrential rains in the area. This is in agreement with the observations made by Trenberth *et al* (2007) that as climate changes, changes are occurring in the amount, intensity, frequency and type of precipitation.




Figure 3: Monthly Rainfall for 2012 and 2013

Increasing frequency and severity of flood do not stem from torrential rainfall alone. It is noted that increasing rate of urbanization in the absent of well-articulated and comprehensive physical planning and land use planning control is one other major factors that causes floods in the area. About 74% of the respondents in the various areas affected by the flood said that their houses plans were not approved by the relevant authority before they were built.

In terms of drainage systems, 64.5% of the respondents said there are no drainages in their communities while 35.5% have drainage systems. This is in-line with the field observations that show inadequate drainage channels in most of the affected areas visited. In areas where drainages exist, these have been blocked by refuse as the communities are in the habit of dumping their wastes in the drainage channels. It was also observed that most of the houses affected were constructed on waterways. About 69% of the houses according to the respondents were built with mud. These mud houses; couple with the sandy nature of the soil cannot easily withstand any flood disasters. About 77.5% of the respondents said that the state government through the Ministry of Environment has been enlightening them on the dangers of farming and building on areas liable to flooding, but people still construct houses (especially in urban LGAs like Katsina and Badagarawa) and others farm in such areas as testified by 69% of the respondents.

Deforestation which is the deliberate destruction of the vegetation as a result of the activities of Man and his domesticated animals (Abaje, 2007), can intensify flooding by affecting the soil structure, reducing infiltration rates and reducing water storage. Evidence has shown that forests and grasses provide an intercepting layer for rainfall, which reduces the rate of overland flow. From the surveyed questionnaires, 72.5% of the respondents use firewood as their source of cooking, 5.5% and 22% use gas and kerosene respectively. With this, it is clear that deforestation is another factor that causes floods in the area. This is evidence from the 78% of the respondents that said they have never planted any tree to replace the ones cut down.

Damages Caused by the Flood Disasters and Human Security

In 2012 and 2013, the ravaging effects of flooding in Nigeria became so drastic that it was seen as a national disaster. Katsina state was among the states that had the bitter experience of the flood. Houses and other public and private properties, infrastructure and facilities were submerged and destroyed, while many residents were displaced (NEMA, 2013). Data collected and analyzed revealed that the damage caused by the flood ranges from loss of lives and livelihood assets to destruction of houses and infrastructural facilities in the areas.

Infrastructures here include bridges, roads, schools, hospitals, markets, electric poles, etc. Analyzing the response of the flood disaster on infrastructural facilities shows the damage on roads were the most severely affected according to 46.5% of the respondents and unfortunately, roads are the largest, fastest-growing and the most heavily used transport infrastructure in the area and Nigeria at large. This is followed by bridges (20%), hospitals/health centers (13.5%), electric poles (12%), schools (5.5%) and markets (2.5%). This means increase maintenance and repair costs, interruption of critical evacuation routes and energy supplies, disruption of economic activity, and degrading of quality of lives among others.

The widespread destruction of infrastructural facilities caused by the floods exerted a huge negative impact on health, power, social and communication sector. The roads that were washed away and collapsed bridges (for example, Charanchi bridge) cut off many communities, making the delivery of health services a difficult task and also making it impossible to get people from other communities to assist flood victims. The health security of the people who have been affected by the floods was jeopardized by the insanitary conditions of those areas, which encouraging the quick spread of water-borne diseases, the spreading of infectious respiratory and skin diseases, spatial and temporal transmission of disease vectors such as malaria, dengue fever, meningitis, and cholera in the area. Besides the weak sanitation and hygiene's conditions in camps which compromised the health security of the people affected, there were increase cases of deaths by drowning rising issues concerning personal and community security of the people affected by the flood disaster.

Since there is interrelationship between infrastructures and other sectors of the economy, a destruction of infrastructures by flooding leads to disruption of socio-economic activities which trigger some vulnerabilities such as malnutrition and food insecurity, poverty, unemployment and economic insecurity among others, which eventually results in developmental problem in the State. For example, the 2012 flood alone affected 10,536 people in the state and a total loss of ₦679, 880, 580:00 (SEMA, 2014).

Conclusions

In assessing the flood disaster in some parts of Katsina state, it was noted that most of the flood occurrences owe their reasons not only to high torrential rainfall, but also improper physical planning, blockage of drainage channels, deforestation and the erection of structures in areas of high risk. It was also noted that flood disaster triggered vulnerability factor as it left many people and communities in precarious conditions, depriving them of most basic goods.

The implications of the increased in rainfall and the frequent occurrences of floods are that models

built on the perceived decreasing rainfall, such as drainages, bridges among others, have to be reviewed. This will lead to increase burden of the recurrent cost of repair/replacement.

The study recommends that: 1) the planning and designing of infrastructures should take into account the nature of rainfall variability in the state; 2) the provision of adequate drainage facilities should be an important segment of all development planning programs in the study area; 3) capacity building to integrate climate change and its image into urban development planning involving local communities, raising public awareness and education on climate change; and 4) the establishment and improvement of early warning systems by Nigerian Meteorological Agency to monitor the occurrence of both floods and droughts in these areas would help in planning of relief measures that would reduce the level of human insecurity in the study area.

References

- Abaje, I.B. & Giwa, P.N. (2010). Flood Risk Assessment and Vulnerability in Kafanchan Town, Jema'a Local Government Area of Kaduna State, Nigeria. *International Journal of Sustainable Development*, 3(1),94-100.
- Abaje, I.B. (2007). *Introduction to Soils and Vegetation*. Kafanchan: Personal Touch Productions.
- Abaje, I.B. Ndabula, C. and Garba, A.H. (2014). Is the Changing Rainfall Patterns of Kano State and its Adverse Impacts an Indication of Climate Change? *European Scientific Journal*, 10 (2), 192-206.
- Abaje, I.B., Ati, O.F. & Iguisi, E.O. (2012a). Changing climatic scenarios and strategies for drought adaptation and mitigation in the Sudano-Sahelian Ecological Zone of Nigeria. In M.A. Iliya & I.M. Dankani (Eds), *Climate Change and Sustainable Development in Nigeria* (pp. 99-121). Ibadan: Crown F. Publishers.
- Abaje, I.B., Ati, O.F. & Iguisi, E.O. (2012b). Recent trends and fluctuations of annual rainfall in the Sudano-Sahelian Ecological Zone of Nigeria: Risks and opportunities. *Journal of Sustainable Society*, 1(2), 44-51.
- Abaje, I.B., Ati, O.F., Iguisi, E.O. & Jidauna, G.G. (2013). Droughts in the Sudano-Sahelian Ecological Zone of Nigeria: Implications for Agriculture and Water Resources Development. *Global Journal of Human Social Science (B): Geography, Geo-Sciences & Environmental*. 13 (2), 1-10.
- Abaje, I.B., Sawa, B.A. & Ati, O.F. (2014). Climate Variability and Change, Impacts and Adaptation Strategies in Dutsin-Ma Local Government Area of Katsina State, Nigeria. *Journal of Geography and Geology*, 6 (2), 103-112.
- Battersby, P. & Siracus, J. (2009). *Globalization and Human Security*. New York: Rowman and Little Field Publishers Inc.
- Box, G.E.P. and D.R. Cox, 1964. An Analysis of Transformation. *Roy. Stat. Soc., B.*, 26: 211-252.
- Federal Republic of Nigeria. (April, 2010). Federal Republic of Nigeria 2006 Population and Housing Census. Priority Table Vol.III. Abuja: National Population Commission
- Katsina State Emergency Management Agency (2014). 2012and 2013 Flood Disasters in Katsina State.
- Nigeria Emergency Management Agency. (2013). Flood Disaster Report 2012, *Daily Trust Newspaper*, 11th May 2013.
- Nigerian Meteorological Agency. (2012). Temperature. *2012 Nigeria Climate Review* (pp 19-24). Nigerian Meteorological Agency (NIMET), Abuja.
- Odekunle, T.O. (2006). Determining Rainy Season Onset and Retreat over Nigeria from Precipitation Amount and Number of Rainy Days. *Theor. Appl. Climatol.* 83, 193–201.
- Odekunle, T.O., Andrew, O. & Aremu, S.O. (2008). Towards a wetter Sudano-Sahelian Ecological Zone in twenty first century Nigeria. *Weather*, 63(3), 66-70.
- Roberts, D. (2008). *Human Security: Global Structure and Violence*. London: Zed Books Ltd.
- Stidd, C.K., 1970. The nth Root Normal Distribution of Precipitation. *Water Resources Research*, 6: 1095-1103.
- Trenberth, K.E., Jones, P.D., Ambenje, P., Bojariu, R., Easterling, D., Klein Tank, A., ...Zhai, P. (2007). Observations: Surface and atmospheric climate change. In Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor M., & Miller, H.L. (Eds), *Climate Change 2007: The Physical Science Basis*, (pp 235-336). Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. United Kingdom and New York, NY: Cambridge University Press.
- United Nations Development Programme. (2009). Linking climate change policies to human development analysis and advocacy. A Guidance Note for Human Development Report Teams.
- Walsh, R.P.D. & Lawler, D.M. (1981). Rainfall seasonality: description, spatial patterns and change through time. *Weather*, 36, 201-208.